

Optical absorption of spin ladders

Christoph Jurecka[†] and Wolfram Brenig^{†,*}

[†]*Institut für Theoretische Physik, Technische Universität Braunschweig, 38106 Braunschweig, Germany*

^{*}*Institute for Theoretical Physics, University of California at Santa Barbara, CA 93106-4030, USA*

(February 1, 2008)

We present a theory of phonon-assisted optical two-magnon absorption in two-leg spin-ladders. Based on the strong intra-rung-coupling limit we show that collective excitations of total spin $S = 0, 1$ and 2 exist outside of the two-magnon continuum. It is demonstrated that the singlet collective state has a clear signature in the optical spectrum.

71.27.+a, 75.10.Jm, 75.50.Ee, 75.30.Et, 78.30.Hv
ITP preprint: NSF-ITP-99121

Recent design of materials with localized spin-1/2 moments arranged in ladder, as well as frustrated and dimerized chain geometries has enormously intensified research on low-dimensional quantum magnetism [1]. Many of the novel materials are likely candidates for a spin-liquid ground state and display a gapped spin-excitation spectrum. On the two-leg ladder the spin-gap is a phenomenon well established by various theoretical approaches, i.e. exact diagonalization [2,3], strong coupling expansion [4,5], density-matrix renormalization group [6], and bosonization [7]. Optical spectroscopy, and in particular Raman scattering, has proven to be a valuable tool to investigate the spin excitations of the new quantum magnets [8]. Apart from Raman scattering, optical absorption has been studied, however, only on the *single-chain* cuprate Sr_2CuO_3 [9,10] where it has provided direct evidence for the gapless two-spinon spectrum of the spin-1/2 Heisenberg chain [11]. In Sr_2CuO_3 absorption is due to phonon assisted two-magnon emission (PME) [12,13]. While several of the new spin-ladder materials, eg. SrCu_2O_3 and $\text{LaCuO}_{2.5}$, may allow for PME as well, corresponding studies are lacking. In this work we propose a simple theory of PME for spin-ladders. We show that *bound states* in the spin-gap which have no analogy in the Heisenberg chain have a substantial effect on the optical absorption spectrum.

We start our discussion with the two-leg spin-ladder Hamiltonian

$$H = \sum_{l,\alpha} [S_{1l}^\alpha S_{2l}^\alpha + \lambda (S_{1l}^\alpha S_{1l+1}^\alpha + S_{2l}^\alpha S_{2l+1}^\alpha)] \quad (1)$$

where $S_{\mu l}^\alpha$ with $\alpha = x, y, z$ is a spin-1/2 operator on site l of leg μ and H is measured in units of $J_\perp$ with $\lambda = J_\parallel/J_\perp$. In this paper we focus on the limit of large intra-rung coupling $\lambda \rightarrow 0$ in which a reformulation of (1) in terms of the rung-spin $S_l^\alpha = S_{1l}^\alpha + S_{2l}^\alpha$ and the operator $T_l^\alpha = S_{1l}^\alpha - S_{2l}^\alpha$ is appropriate

$$H = \sum_l \left[-\frac{3}{4} + \frac{1}{2} S_l(S_l + 1) \right] + \frac{\lambda}{2} \sum_{l,\alpha} T_l^\alpha T_{l+1}^\alpha$$

$$+ \frac{\lambda}{2} \sum_{l,\alpha} S_l^\alpha S_{l+1}^\alpha = H_1 + H_2 + H_3 \quad (2)$$

where $S_l = 0(1)$ for a rung singlet(triplet). The rung-spin eigenbasis is given by $|s\rangle = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/\sqrt{2}$, $|t_x\rangle = -(|\uparrow\uparrow\rangle - |\downarrow\downarrow\rangle)/\sqrt{2}$, $|t_y\rangle = i(|\uparrow\uparrow\rangle + |\downarrow\downarrow\rangle)/\sqrt{2}$, and $|t_z\rangle = (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2}$ where the first(second) entry in the kets refers to a site on leg '1(2)' of the ladder. As usual $S^\alpha|s\rangle = 0$ and $S^\alpha|t_\beta\rangle = i\varepsilon_{\alpha\beta\gamma}|t_\gamma\rangle$ with the Levi-Civita symbol $\varepsilon_{\alpha\beta\gamma}$. The action of T^α on the rung basis is given by $T^\alpha|s\rangle = |t_\alpha\rangle$ and $T^\alpha|t_\beta\rangle = \delta_{\alpha\beta}|s\rangle$.

For vanishing inter-rung coupling the ground state of (2) is a pure rung-singlet product-state $|\rangle = \otimes_l |s_l\rangle$. The excited states are products of singlets and triplets $|\{t_{m\alpha}\}\rangle = \otimes_{\{l\}} |s_l\rangle \otimes_{\{m\alpha\}} |t_{m\alpha}\rangle$ with an excitation energy given by the number of triplets $N[\{t_{m\alpha}\}]$. At finite λ the action of the ladder-Hamiltonian on these product states can be read off easily from (2): (i) H_2 creates(destroys) pairs of nearest-neighbor (NN) triplets of equal α -index, (ii) given a pair of NN sites in a relative state of one singlet and one triplet H_2 generates NN hopping of the triplet via exchange of the singlet and triplet, and finally (iii) H_3 induces NN interactions between triplets. Contributions (ii) and (iii) define a three-flavored hard-core Bose system with NN hopping as well as NN interactions. These contributions renormalize the spectrum to $O(\lambda)$. Process (i) does not conserve the triplet number and changes the spectrum only to $O(\lambda^2)$ at $\lambda \ll 1$. The essential physics of the bound states and optical absorption is independent of $O(\lambda^2)$ -terms. They will be neglected hereafter. Based on this simplification it is convenient to split the Hamiltonian into a *bare* 'kinetic' part $H_1 + H_2$ and a two-particle interaction H_3 . The *bare* one-triplet eigenstates $|k\alpha\rangle$ of momentum k are

$$|k\alpha\rangle = \frac{1}{\sqrt{N}} \sum_l e^{ikl} |t_{l\alpha}\rangle \quad (3)$$

$$\varepsilon_k = \langle k\alpha | H_1 + H_2 | k\alpha \rangle - \langle |H| \rangle = 1 + \lambda \cos(k) \quad (4)$$

where $\langle k'\alpha | k\beta \rangle = \delta_{\alpha\beta} \delta_{kk'}$. This agrees with seminal work on spin-ladders [2-4]. Hereafter the ground state energy $\langle |H| \rangle$ is set to zero. The *bare* two-triplet eigenstates in the singlet sector must incorporate the symmetry $|t_{l\alpha} t_{m\beta}\rangle = |t_{m\beta} t_{l\alpha}\rangle$ as well as the hardcore constraint $|t_{l\alpha} t_{l\beta}\rangle = 0$

$$|kq, S\rangle = \frac{1}{\sqrt{N(N-1)}} \sum_{l,m} [e^{ik(l+m)/2} \text{sgn}(l-m) \sin(q(l-m)) |lm, S\rangle] \quad (5)$$

$$\varepsilon_{kq} = 2[1 + \lambda \cos(k/2) \cos(q)] \quad (6)$$

where $|lm, S\rangle = \sum_{\alpha} |t_{l\alpha} t_{m\alpha}\rangle / \sqrt{3}$ refers to the singlet combination of two triplets in real space and ε_{kq} is the bare two-particle kinetic energy. $k(2q)$ is the total(relative) momentum, $\langle k'q' | kq \rangle = \delta_{kk'} \delta_{qq'}$ and $k \in]-\pi, \pi]$, $q \in]0, \pi]$. (Anti)periodic boundary conditions apply to $k(q)$ and $|k, q\rangle = -|k, -q\rangle$. A discussion of the remaining two-particle states in the triplet and quintet sector is deferred to appendix A.

After these preliminaries we are in a position to evaluate the optical absorption due to *phonon-assisted two-magnon emission* (PME). This absorption results from a magnetoelastic excitation induced by the incoming photon-field and has been considered first in the planar geometry of the CuO_2 square-lattice of the HT_C materials [12,13]. Its microscopic justification can be applied with little change to the case of spin-ladders with corner-sharing Cu-O structure. For the effective coupling one obtains

$$H_{LS} = -E(t) \sum_{l,\alpha,\mu} [p_I u_{\mu Ol} + p_A (2u_{\mu Ol} - u_{\mu Ll} - u_{\mu Rl})] S_{\mu l}^{\alpha} S_{\mu l+1}^{\alpha} \quad (7)$$

$E(t)$ is the time-dependent electric field which is polarized along the ladder. $u_{\mu O(L,R)l}$ denote displacement coordinates of the oxygen site *on* the μ -th leg of the ladder and refer to the oxygen sites in cell $O(\text{right}) = l$, $L(\text{left}) = l - 1$, and $R(\text{right}) = l + 1$. The effective charges p_I and p_A can be derived from an expansion up to first order in u_l and $E(t)$ of the Cu-O-Cu superexchange coupling. Expressions for p_I and p_A can be found in [12,13]. For cuprates which are strongly covalent materials and for $E(t)$ polarized parallel to the ladder, $|p_A| \gg |p_I|$ is a reasonable approximation [13]. Although the specific form of (7) depends on the corner-sharing Cu-O structure it is conceivable that a similar PME is also possible in other spin-ladder compounds. Introducing the Fourier transformed phonon-operators $u_{k\mu} = (a_{k\mu} + a_{-k\mu}^{\dagger}) / \sqrt{2M\omega_k}$ with the phonon dispersion ω_k and the reduced O mass M and replacing $S_{\mu l}^{\alpha}$ with the bond variables S_l^{α} and T_l^{α} we get

$$H_{LS} = E(t) \frac{1}{\sqrt{2}} \sum_{k,\mu} (a_{k\mu} + a_{-k\mu}^{\dagger}) P_k \quad (8)$$

$$P_k = -\frac{1}{4\sqrt{NM}\omega_k} \sum_{l,\alpha} e^{ikl} [p_I + 2p_A(1 - \cos(k))] T_l^{\alpha} T_{l+1}^{\alpha} \\ := -\frac{1}{\sqrt{N}} \sum_{l,\alpha} p_k e^{ikl} T_l^{\alpha} T_{l+1}^{\alpha} \quad (9)$$

where terms proportional to $S_l^{\alpha} S_{l+1}^{\alpha}$ and $S_l^{\alpha} T_{l+1}^{\alpha}$ have been dropped since they do not act on a pure singlet product-state. Photon momentum conservation has been neglected since the wave length of light is large compared to the lattice spacing.




FIG. 1. Dispersion of the $S = 0, 1$, and 2 bound states relative to the bare two-triplet continuum.

The zero-temperature absorption coefficient $I(\omega)$ at frequency ω , normalized to the electric field intensity $|E|^2$, is obtained from Fermi's golden rule. To simplify technical matters we assume only optical phonons of frequency $\omega_k = \omega_0$ to be involved in the absorption process. Since our model discards any intrinsic magnetoelastic coupling the phonons represent a mere 'momentum bath' allowing for arbitrary recoil of the two triplets [14]. Tracing over the phonons leads to

$$I(\omega) = 2\pi \sum_{f,k} |\langle fk | P_k | 0 \rangle|^2 \delta(\omega - E_{fk} - \omega_0) \quad (10)$$

$$= -2 \text{Im} \sum_{k,q,q'} [\langle 0 | P_k^{\dagger} | kq, S \rangle \langle kq, S | \frac{1}{z - H} | kq', S \rangle \langle kq', S | P_k | 0 \rangle] \quad (11)$$

where $z = \omega + i0^+ - \omega_0$ and for the remainder of this work we redefine $\omega - \omega_0 \rightarrow \omega$. While $|fk\rangle$ are the *dressed* two-particle eigenstates of the ladder of total momentum k , spin zero and energy E_{fk} , $|kq, S\rangle$ are the bare states (5).

Next we have to evaluate the resolvent (11). Since $|kq, S\rangle$ contains only two triplets and, as noted previously, we neglect the coupling between sectors of different triplet number this task is equivalent to solving an interacting two-particle problem which can be done exactly. To this end we observe that

$$\langle kq, S | H_3 | k'q', S \rangle = -\frac{4}{N-1} \delta_{kk'} \lambda \sin(q) \sin(q') \quad (12)$$

$$\langle k'q, S | P_k | 0 \rangle = -\sqrt{\frac{12}{N-1}} p_k e^{-ik/2} \sin(q) \delta_{kk'} \quad (13)$$

This can be combined with (11) to sum the T-Matrix series for $\langle kq, S | 1/(z - H) | kq', S \rangle$ yielding

$$I(\omega) = -24 \text{Im} \sum_k \frac{p_k^2}{G^{-1}(k, \omega + i0^+) + 4\lambda} \quad (14)$$




FIG. 2. Optical absorption spectrum for $p_I = 0$. Inset: identical spectrum, y-axis stretched by 100.

where the bare two particle resolvent $G(k, z)$ is given by

$$G(k, z) = \frac{1}{4\pi} \int_{-\pi}^{\pi} dq \frac{\sin^2(q)}{z - (2 + 2\lambda \cos(k/2) \cos(q))}$$

$$= \frac{\text{sign}[\text{Re}(a)]\sqrt{a^2 - 1} - a}{4\lambda \cos(k/2)} \quad (15)$$

with $a = (2 - z)/[2\lambda \cos(k/2)]$. The zeros at $\omega = E_k^S$ and momentum k of the denominator on the r.h.s. of (14)

$$1 + 4\lambda G(k, E_k^S) = 0 \quad (16)$$

correspond to (anti)bound states in the two-triplet spin-singlet sector. Since both, the boundaries of the two-particle continuum (6) as well as the solutions of (16) are functions of k only if energies ω are rescaled in terms of the quantity $(\omega - 2)/\lambda$, the k -space structure of the rescaled two-particle spectrum is *independent* of λ within our approximation. In particular, a singlet bound-state exists at all k in the Brillouin zone. At $k = \pi$ its binding energy is largest with $E_k^S = 2 - \lambda$ while at $k = 0$ the binding energy is zero. This is shown in fig. 1, which depicts the bound-state dispersion along with the two-particle continuum. Also shown are triplet ($S = 1$) and quintet ($S = 2$) (anti)bound-states, which are optically inactive. These states are discussed in appendix A. Recently, similar collective states have been predicted using bond-boson techniques [15–17].

From (13,14,15) we can obtain the absorption by numerical integration. The rescaled intensity $\lambda I(\tilde{\omega})/p_A$ is a function of $\tilde{\omega} = (\omega - 2)/\lambda$ and p_I/p_A only and does not depend on λ . Figure 2 depicts the absorption spectrum for $p_I = 0$. We note that our choice of the frequency variable $\tilde{\omega}$ implies that zero incoming photon energy corresponds to the point $-2/\lambda$ on the x-axis of fig. 2. For any finite spin gap, i.e. $\lambda < 1$, this point is




FIG. 3. Optical absorption spectrum for $p_I = -2p_A$.

off the range plotted which refers only to the frequency window $-2 \leq \tilde{\omega} \leq 2$ in which absorption occurs. Figure 2 demonstrates that the bound state has a profound impact on the absorption spectrum which comprises almost completely of a structure due to the integrated density of states of the bound state. In particular there is a van-Hove singularity at $\omega = 2 - \lambda$ which results from the maximum in the bound-state dispersion at $k = \pi$. Only in the special case of $p_I = -4p_A$ this van-Hove singularity disappears. The inset in fig. 2 focuses on the remaining spectral weight. It has a maximum at the center of the two-triplet continuum, however its weight is very small as compared to the contribution from the bound states. At finite values of p_I a step-like feature appears at the spin-gap and for $p_I < 0$ a dip occurs due to interference in the effective charge p_k . At present we believe that such features are of academic interest only. Yet, for completeness a typical spectrum displaying the latter effects is shown in fig. 3.

The preceding discussion does not rule out an interpretation of van-Hove like structures in an experimentally determined absorption spectrum solely in terms the momentum integrated *bare* two-triplet continuum rather than in terms of bound states. However, taking this point of view and using that $|p_I/p_A| \ll 1$, which is most likely the case, the absorption spectrum would rather be symmetric with respect to the center of gravity of the measured spectrum. Finally we note, that in case of a finite dispersion of the phonon additional structures may appear which are not contained in our present analysis.

In conclusion we propose that singlet bound-states have a dramatic impact on the optical absorption of dimerized spin-ladder systems which allow for phonon-assisted multi-magnon absorption of the Lorenzana-Sawatzky type. We hope that our work may stimulate further experimental investigation of this issue.

One of us (W.B.) acknowledges the kind hospitality of the ITP at UC Santa Barbara. This research was sup-

ported in part by the Deutsche Forschungsgemeinschaft under Grant No. BR 1084/1-1 and the National Science Foundation under Grant No. PHY94-07194.

APPENDIX A: S=1,2 STATES

The triplet and quintet two-particle states in real space correspond to the combinations $|lm, \alpha\rangle = \sum_{\beta, \gamma} \varepsilon_{\alpha\beta\gamma} |t_{l\beta} t_{m\gamma}\rangle / \sqrt{2}$ and $|lm, \alpha\beta\rangle = \sqrt{3/8}(|t_{l\alpha} t_{m\beta}\rangle + |t_{l\beta} t_{m\alpha}\rangle - \delta_{\alpha\beta}(2/3) \sum_{\gamma} |t_{l\gamma} t_{m\gamma}\rangle)$ respectively. Due to spin rotational invariance we may focus on the total- $S^z = 0$ components only, i.e. $|lm, P\rangle = |lm, z\rangle$ (triplet) and $|lm, D\rangle = |lm, zz\rangle$ (quintet). The bare two-particle eigenstates are

$$|kq, P\rangle = \frac{1}{N} \sum_{l, m} e^{i[k(l+m)/2 + q(l-m)]} |lm, P\rangle \quad (\text{A1})$$

$$|kq, D\rangle = \frac{1}{\sqrt{N(N-1)}} \sum_{l, m} [e^{ik(l+m)/2} \text{sgn}(l-m) \sin(q(l-m)) |lm, D\rangle] \quad (\text{A2})$$

where identical properties regarding the momenta as in (5) apply and the bare two-particle energies of the states $|kq, P\rangle$ and $|kq, D\rangle$ are identical to ε_{kq} of (5). Note that both, P - and D -states conform with the hardcore constraint $|t_{l\alpha} t_{l\beta}\rangle = 0$. From the identity $(\mathbf{S}_l + \mathbf{S}_m)^2/2 - 2 = \mathbf{S}_l \cdot \mathbf{S}_m$ and from (12) it follows directly that

$$\langle kq, X | H_3 | k'q', X \rangle = -\frac{c_X}{N-1} \delta_{kk'} \lambda \sin(q) \sin(q') \quad (\text{A3})$$

with $c_X = 4, 2$, and -2 for $X = S, P$, and D respectively. Therefore the energies E_k^X of the (anti)bound states in all spin channels are obtained from the zeros of the single equation

$$1 + c_X \lambda G(k, E_k^X) = 0 \quad (\text{A4})$$

with $G(k, z)$ as in (15) and c_X set according to the total spin. For $X = P(D)$ a(n) (anti)bound state exists for $k \geq 2\pi/3$ for all values of λ with $E_\pi^{P(D)} = 2 - (+)\lambda/2$. In fig. 1 the dispersion of these (anti)bound states relative to the two-triplet continuum is depicted.

- [5] J. Oitmaa, R.R.P. Singh, Z. Weihong. Phys. Rev. B **54**, 1009 (1996)
- [6] S.R. White, R.M. Noack, and D.J. Scalapino, Phys. Rev. Lett. **73**, 886 (1994)
- [7] D.G. Shelton, A.A. Nersesyan, and A.M. Tsvelik, Phys. Rev. B **53**, 8521 (1996)
- [8] P. Lemmens, M. Fischer, M. Grove, P.H.M. v. Loosdrecht, G. Els, E. Sherman, C. Pinettes, G. Güntherodt, cond-mat/9910074, *and refs. therein*
- [9] H. Suzuura, H. Yasuhara, A. Furusaki, N. Nagaosa, and Y. Tokura, Phys. Rev. Lett. **76**, 2579 (1996)
- [10] J. Lorenzana and R. Eder, Phys. Rev. B **55**, R3358 (1997)
- [11] A.H. Bougourzi, M. Karbach, and G. Müller, Phys. Rev. B **57**, 11429 (1997)
- [12] J. Lorenzana and G.A. Sawatzky, Phys. Rev. Lett. **74**, 1867 (1995)
- [13] J. Lorenzana and G.A. Sawatzky, Phys. Rev. B **74**, 9576 (1995)
- [14] This is different from two-magnon Raman scattering which probes zero total two-magnon momentum only and may therefore be less well suited to observe $S = 0$ bound states with dominant weight at the zone boundary.
- [15] K. Damle and S. Sachdev, Phys. Rev. B **57**, 8307 (1998)
- [16] O.P. Sushkov and V.N. Kotov Phys. Rev. Lett. **81**, 1941 (1998)
- [17] While in the present work two triplets never occupy the same site by construction of (5,A1,A2) a fictitious hardcore potential has to be dealt with in refs. [15,16] in order to remove unphysical states containing double-occupancies from the Hilbert space.

-
- [1] For a recent review on ladder materials see: E. Dagotto, cond-mat/9908250, *and refs. therein*
 - [2] T. Barnes, E. Dagotto, J. Riera, and E. Swanson, Phys. Rev B **47**, 3196 (1993)
 - [3] T. Barnes and J. Riera, Phys. Rev B **50**, 6817 (1994)
 - [4] M. Reigrotzki, H. Tsunetsugu, and T.M. Rice, J. Phys. Condens. Matter **6**, 9235 (1994)